

Evolution of medical treatment for endometriosis: back to the roots?

Fritz Wieser^{1,2,6}, Misha Cohen³, Andrew Gaedert⁴, Jie Yu², Carla Burks-Wicks⁵, Sarah L. Berga² and Robert N. Taylor²

¹Division of Gynecological Endocrinology and Reproductive Medicine, University of Vienna, 1090 Vienna, Austria; ²Department of Gynecology and Obstetrics, Emory University School of Medicine, Woodruff Memorial Building, 1639 Pierce Drive, Room 4217, Atlanta, GA 30322, USA; ³UCSF Institute for Health and Aging, San Francisco, CA, USA; ⁴Health Concerns, Oakland, CA, USA; ⁵Department of Obstetrics and Gynecology, UCSF, San Francisco, CA, USA

⁶Correspondence address. Tel: +1-404-727-9823; Fax: +1-404-727-8609; E-mail: fwieser@emory.edu

Experimental evidence is accumulating to suggest that medicinal botanicals have anti-inflammatory and pain-alleviating properties and hold promise for treatment of endometriosis. Herein, we present a systematic review of clinical and experimental data on the use of medicinal herbs in the treatment of endometriosis. Although there is a general lack of evidence from clinical studies on the potential efficacy of medicinal herbs for the treatment of endometriosis-associated symptoms, our review highlights the anti-inflammatory and pain-alleviating mechanisms of action of herbal remedies. Medicinal herbs and their active components exhibit cytokine-suppressive, COX-2-inhibiting, antioxidant, sedative and pain-alleviating properties. Each of these mechanisms of action would be predicted to have salutary effects in endometriosis. Better understanding of the mechanisms of action, toxicity and herb–herb and herb–drug interactions permits the optimization of design and execution of complementary alternative medicine trials for endometriosis-associated pain. A potential benefit of herbal therapy is the likelihood of synergistic interactions within individual or combinations of plants. In this sense, phytotherapies may be analogous to nutraceuticals or whole food nutrition. We encourage the development of herbal analogues and establishment of special, simplified registration procedures for certain medicinal products, particularly herbal derivatives with a long tradition of safe use.

Keywords: endometriosis; herbs; botanicals; inflammation; CAM

Introduction

The successful treatment of endometriosis-associated symptoms including dysmenorrhoea, dyspareunia and chronic non-menstrual pain typically requires surgical as well medical intervention (Kennedy *et al.*, 2005). The use of both general modalities for endometriosis-associated pain has been recently reviewed (Kennedy *et al.*, 2005; Crosignani *et al.*, 2006). The clinical prevalence of endometriosis symptomatology is very high. In a survey of reproductive-aged women in Canada, 60% met the criteria for primary dysmenorrhea (Burnett *et al.*, 2005). In adult women with endometriosis, the prevalence rate of dysmenorrhoea was up to 76% (Kuohung *et al.*, 2002), and in adolescents with endometriosis, it was up to 94% (Reese *et al.*, 1996). This disease constitutes a public health dilemma of major proportion. Although medical therapies are not curative *per se*, they are nonetheless a mainstay of pain symptom suppression among women with endometriosis.

Initially high-dose diethylstilbestrol and combinations of potent estrogens and progestagens were used to treat endometriosis (Hurxthal and Smith, 1952), but this approach was subsequently

replaced by progestagens alone (Kistner, 1958). In 1958, the clinical observation of an apparent resolution of symptoms during pregnancy gave rise to the concept of treating patients with pseudo-pregnancy hormone regimens (Kistner, 1958). Different forms of progestagens and anti-progestagens (including dihydroprogesterone, medroxyprogesterone acetate (MPA), gestrinone and mifepristone (RU 486) afforded some improvement in pelvic pain; however, these treatments are ineffective in a subset of women with endometriosis and carry several untoward side effects (Vercellini *et al.*, 1997). In 1973, danazol, a isoxazole derivative of 17-alpha testosterone, was introduced for the treatment of endometriosis-associated pain (Friedlander, 1973). Although danazol is effective, extended use is limited by androgenic and metabolic side effects (Selak *et al.*, 2001). A decade later, in 1982, gonadotrophin releasing hormone agonists (GnRH-a) were first described as an alternative treatment for endometriosis (Lemay and Quesnel, 1982). Although GnRH-a can be used safely with combined estrogen and/or progestagen add-back therapy for up to 2 years, long-term use is constrained

by hypoestrogenic side effects (Corson and Bolognese, 1978), especially in adolescents.

Since 1978, non-hormonal treatment regimens for endometriosis-associated pain, including a variety of non-steroidal anti-inflammatory drugs (NSAIDs), have been promoted (Corson and Bolognese, 1978). Objective evidence on the use of NSAIDs in endometriosis-associated pain is sparse and inconclusive (Allen *et al.*, 2005). Despite the poor quality of evidence, NSAIDs are typically used as first-line drugs in the treatment of endometriosis associated-pain, because they are felt to have fewer limitations (Allen *et al.*, 2005).

Much effort is spent on the development and promotion of new drug treatments with the goal of achieving higher efficacy, fewer side effects and the option of long-term treatment, especially in women with severe endometriosis. These agents include thiazolidinediones (Lebovic *et al.*, 2004), selective progesterone receptor modulators (SPRMs) (Chwalisz *et al.*, 2005), aromatase inhibitors (Amsterdam *et al.*, 2005), cyclooxygenase (COX)-2 selective NSAIDs (Cobellis *et al.*, 2004), recombinant human TNF-alpha binding proteins (Barrier *et al.*, 2004), anti-VEGF therapy (Nap *et al.*, 2005), MMP-inhibitors (Mori *et al.*, 2001) and interferon-alpha-2b (Badawy *et al.*, 2001). Evidence from pre-clinical trials has suggested beneficial effects of these drugs that may be conferred by anti-proliferative, anti-inflammatory or anti-angiogenic mechanisms. However, despite these many therapeutic options, efficient long-term regimens for the treatment of endometriosis-associated symptoms are desperately needed.

In recent years, medicinal herbs and other botanical products have become popular for management of symptoms of several gynaecologic disorders (Eisenberg *et al.*, 1998; Anderson and Johnson, 2005; Comar and Kirby, 2005; Tindle *et al.*, 2005) including endometriosis-associated symptoms (Cox *et al.*, 2003; Fugh-Berman and Kronenberg, 2003). Evidence for the potential efficacy of medicinal herbs in the treatment of endometriosis-associated symptoms has been reported in the literature and is the focus of this review.

Background

Traditional medicine practices refer to health approaches, knowledge and beliefs incorporating natural plant, animal and mineral-based medicines, spiritual therapies, manual techniques and exercises (Chen and Chen, 2004; Huang, 1998). These can be applied individually or in combination to treat and prevent illnesses or maintain well-being. In industrialized countries, adaptations of traditional medicine are termed complementary and alternative medicine (CAM). CAM is popular in all regions of the developing world and its use is rapidly spreading in industrialized countries. In Europe and North America, over 50% of the population have used CAM at least once (Tindle *et al.*, 2005).

Historically, Chinese culture has relied heavily on herbal treatment of many illnesses. Traditional herbal preparations still account for 30–50% of the total medicinal consumption in China. Written records document the use of Chinese herbal medicine over 3000 years ago. In Chinese medicine, endometriosis is called Neiyi and is considered a ‘Blood stasis syndrome’ resulting in the formation of endometriotic lesions (Flaws, 1989; Maciocia, 1997). Chinese herbal formulae designed for endometriosis therapy are targeted to resolve

blood stasis. Despite centuries of use abroad, medicinal herbal treatments for endometriosis-associated symptoms were only introduced in the USA in the mid-1980s. Several medicinal herbs that historically were prescribed for treatment of endometriosis-associated symptoms are still in use today (Chen and Chen, 2004; Huang, 1998) (Table 1). Each of the herbs described is likely composed of several active components with anti-inflammatory, anti-proliferative and pain-alleviating properties (Table 2). Thus, the potential for multiple synergistic interactions is enormous.

Clinical evidence on the efficacy, toxicity and herb–drug interactions of medicinal herbs and herbal combinations used in the treatment of endometriosis

We searched Medline (1969 to November 2006), EMBASE (1984 to April 2006) and hand searched several prominent journals published in China. Clinical evidence on the efficacy of herbal combinations in treatment of endometriosis is almost exclusively published in the Chinese scientific literature. As Chinese medicine traditionally uses mixtures of medicinal plants, synergistic as well as interfering effects may occur. The concept of synergistic interaction refers to the possibility that when two (or more) active substances are given concurrently, the substances may interact enhancing the effect of the other and at lower doses. Alternatively, multiple compounds could result in decreased efficacy. These possible interactions between two or more drugs/herbs are classified into pharmacokinetic and pharmacodynamic effects (Harrison and Fauci, 1998). Examples of medicinal herbal combinations that have been used to treat endometriosis-associated symptoms include: dan’e recipe, Gexia Zhuyu Tang, Keishi-bukuryo-gan (KBG), Neiyi recipe #1, Neiyi recipe #2, Neiyixiao recipe, Shixiao San (Sudden Smile Powder), Shaofu Zhuyu Tang, Shixiao Guijie Tang, Taoren Chenqi Tang, Tongjin San, Tuo Mo Tang, Tao He Cheng Tang, Xuefu Zhuyu Tang, Xiao Yao San, Yi Wei Zhu Yu Fang and Yiweining (YWN) (Shao, 1980; Cai, 1982; Dai, 1982; Cao, 1983; Liu *et al.*, 1983; Lin *et al.*, 1988; Zhuang and Xia, 1990; Jin, 1991; Li, 1991; Wang *et al.*, 1991; Qu, 1992; Yu *et al.*, 1993; Liu, 1994; Hu and Li, 1995; Liu *et al.*, 1998; Wang *et al.*, 1998; Cai *et al.*, 1999; Yang *et al.*, 2006a) (Table 3).

Unfortunately, the standards of evidence-based principles are sporadically applied in published clinical studies in this area. Published trials of medicinal herbs concerning endometriosis in the Chinese scientific journals of complementary medicine have generally not been conducted according to the guidelines of evidence-based medicine in Western scientific journals. One of the major difficulties in studying the clinical effects of herbal combinations in respect to evidence-based standards is that the composition of herbal formulae is individualized for each patient according to the different syndromes of endometriosis (‘blood stasis syndrome’). Only one randomized controlled trial on the potential effects of medicinal herbs in endometriosis was identified after a thorough search in the English literature (Table 3) (Yang *et al.*, 2006a). This study demonstrated that YWN (consisting of Chinese angelica, corydalis, curcuma, persica, red peony, safflower, salvia root and tortoise shell) was safe and its efficacy was similar to gestrinone in the prevention of post-operative recurrences of endometriosis. Other trials have been published only as

Table 1: Medicinal herbs and natural compounds used in the treatment of endometriosis and their biochemical constituents

English name	Chinese name	Main constituents
Bupleurum	Chai Hu	Triterpenoids (saikosaponin A–E, saikogenin F), essential oils, coumarin, flavone, carbohydrates
Chinese angelica	Dang Gui	Essential oils (ligustilide, n-butylphtalide), ferulic acid, scopletin
Cattail pollen (Typha)	Pu Huang	Isorhamnetin, pentacosane, alpha-sitosterol, arachidonic acid, palmitic acid, quercetin, sporopellin
Cinnamon twigs	Gui zhi	Essential oils (cinnamic-aldehyde, cinnamic acid, cinneylanine); coumarin
Cnidium fruit	Chuang Xiong	Alkaloids (tetramethylpyrazine); organic acids (ferulic acid, folic acid linoleic acid); essential oils; ligustilide
Corydalis	Yan Hu Suo	D-corydaline, corydalis L, dl-tetrahydropal-matine, coptisine, columbamine
Curcuma zedoria	Yu Jin	Essential oils, curcumin, d-camphene, demethoxy-curcumin, turmerone, starch, lipids, carvone
Cyperus	Xiang Fu	Cyperene, cyperol, cyperol, isocyperol, cymene, limonene, camphene, cyperol, alpha-rotunol
Dahurian angelica	Bai Zhi	Coumarins; byakangelicol, cnidilin, scopoletin, phelopterin, xanthotoxin; essential oil
Frankincense	Ru Xiang	Triterpene acids, alpha-boswellic acids, beta-boswellic acids, basorin, alpha-phellandrene, pinene, arabic acid
Licorice root	Gan Cao	Triterpenoids (glycyrrhizin, glycyrrhetic acid), flavonoids (liguridine, liquiritigenin, uralene)
Myrrh	Mo Yao	Terpenoids; heerabomyrrholic acid, commiphoric acid, cinnamic aldehyde, resin, cuminaldehyde
Persica	Tao Ren	Essential oils (gibberellin A5); emulsin, allantoinase, lipids, vitamin B1
Poria	Fu Ling	Pachymose; pachyman, pachymeran, poriacic acid A, B and C, tumulosic acid
Red peony root	Chi Shao	Paeoniflorin, oxypaeoniflorin, lactiflorin, benzoylpaeoniflorin, daucosterol, d-catechin
Rhubarb	Da Huang	Senoside (A–F), emodine, rhein, physcion, physcion monoglucoside, emodin monoglucoside
Salvia root	Dan Shen	Tanshinone (I, II _A , II _B), militirone, tanshinol A, B, C, salviol, vitamin E
Scutellaria	Huang Qin	Baicalin, beicalein, wogonin, wogonoside, chrysin, tenaxin II, oroxylin A, skullcapflavone I, II
Sparganium	San Leng	Essential oils, starch
Tortoise shell ^a	Bie Jia	Gelatin, collagen, colloid, vitamin D, keratin
Tumeric	Jiang Huang	Curcumin, demethoxycurcumin, turmerone, arturmerone, curcuminoids, phellandrene, zingiberene
White Peony root	Bai Shao	Paeoniflorin, lactiflorin, paeonin, oxypaeoniflorin, hydroxypaeoniflorin, daucosterol

^aTurtle/tortoise products have difficulty passing US customs with valid Cites certificates. We are not endorsing the use of animal-derived medicinals.

personal communications or as clinical observations in the English literature. At present, a randomized controlled trial is in progress in the USA evaluating the use of traditional Chinese medicine (TCM) at the Women's Health Research Unit of the Oregon Health Sciences University and the Oregon College of Oriental Medicine. The protocol (NCT00034047) is funded by the National Center for CAM (NCCAM). The study tests the hypothesis that TCM (acupuncture and medicinal herbs) will reduce endometriosis-related pelvic pain as effectively as nafarelin therapy without causing the pseudo-menopausal side effects that accompany GnRH-a therapy. Women were randomly assigned to receive either 12 weekly treatments of TCM or 12 weeks of the FDA-approved GnRH-a treatment. This will likely prove to be a landmark study in alternative endometriosis therapy.

Despite promising knowledge supporting the potential efficacy of herbal treatments in endometriosis and reports on the efficacy of medicinal herbs in related conditions (e.g. over 80% of pain relief, Table 3), the clinical effects of medicinal herbs on endometriosis remain unclear. Nevertheless, postulated experimental mechanisms of medicinal herbs include demonstrated cytokine suppression, COX-2 inhibition and antioxidant and antinociceptive activities. Treatment of endometriosis-associated symptoms with non-hormonal regimens is a particularly interesting treatment option for adolescent women with endometriosis. Hormonal anti-endometriosis therapies such as GnRH-a, progestins and oral contraceptives bear potential serious side effects, which limit their use in adolescents. As the efficacy of NSAIDs in women with endometriosis remains unclear, non-hormonal treatment of symptoms

with medicinal herbs or herbal components may represent an innovative alternative for adolescents with endometriosis. Moreover, since herbal medicines may exert synergistic effects, these combinations may carry potential benefits of higher efficacy while minimizing toxicity.

Toxicity and herb–drug interactions

There is a potential for toxicity and untoward herb–drug interactions using medicinal herbs in the treatment of endometriosis (Table 4). Evidence on toxicity and herb–drug interactions of medicinal herbs used in the treatment of endometriosis is limited (Hoskins, 1984; Moing *et al.*, 1987; Perry *et al.*, 1990; Takasuna *et al.*, 1995; Kuboniwa *et al.*, 1999; Page and Lawrence, 1999; Ishihara *et al.*, 2000; Li *et al.*, 2001; Amato *et al.*, 2002; Chainani-Wu, 2003; Elinav and Chajek-Shaul, 2003; Ikegami *et al.*, 2003; Joshi *et al.*, 2003; Wong and Chan, 2003; Chen and Chen, 2004; Wojcikowski *et al.*, 2004a,b; Zhou *et al.*, 2004; de Boer *et al.*, 2005; Hu *et al.*, 2005; Kelly *et al.*, 2005; Ammon, 2006; Lao *et al.*, 2006; Xie *et al.*, 2006) (Table 4). Toxicity and untoward herb–drug interactions depend on factors associated with drugs (dose, dose regimen and therapeutic range) and as well as the consumer (age, genetic polymorphism, gender and pathological conditions). Toxicity and herb–drug interactions are unlikely to be evaluated because of the lack of current federal regulations in some industrialized countries, including the USA, and the paucity of research funding in this area. Established monitoring of adverse events caused by herb–drug

Table 2: Medicinal herbs and natural compounds used in the treatment of endometriosis and their anti-inflammatory effects

Herbs	Antiproliferative	Antinociceptive, sedative	Anti-inflammatory action		
			Antioxidant	Suppression	
				COX-2	Cytokines
Bupleurum	+			+	+
Chinese angelica	+	+	+	+	
Dahurian angelica root	+	+		+	+
Cattail pollen				+	
Cinnamon twigs		+		+	
Cnidium fruit	+	+			
Corydalis		+		+	
Curcuma (Turmeric)	+	+	+	+	+
Cyperus	+		+		
Frankincense	+	+		+	+
Licorice root	+	+		+	+
Myrrh		+	+	+	
Persica				+	
Poria	+	+		+	
Red peony root			+		
Rhubarb	+	+		+	+
Salvia root		+	+		+
Scutellaria	+			+	
Sparganium		+		+	
Tortoise shell ^a					+
White peony root				+	

^aTurtle/tortoise products have difficulty passing US customs with valid Cites certificates.

interactions and increased financial support for studies investigating herb–drug interference would provide crucial information regarding public safety.

Effects of medicinal herbs, their active components and herbal combinations on cytokine expression

Suppression of the NF-Kappa B pathway and pro-inflammatory cytokines has been recognized as a major mechanism of conventional drug treatments for endometriosis; progestogens, GnRH-a, danazol and NSAIDs are known to demonstrate cytokine-suppressive activity (Boucher *et al.*, 2000; Zhao *et al.*, 2002; Wieser *et al.*, 2005c; Yang *et al.*, 2006b). Cytokines and chemokines such as tumour necrosis factor (TNF)-alpha, interleukin (IL)-1, IL-6, IL-8, monocyte chemotactic protein (MCP)-1 and regulated on activation, normal T cell expressed and secreted (RANTES) contribute to the pathogenesis of endometriosis by enhancing attachment, angiogenesis and/or proliferation of ectopic endometrial tissues in the pelvis (Bergqvist *et al.*, 2001; Lebovic *et al.*, 2001; Maas *et al.*, 2001).

Curcuma zeodaria is used in many anti-endometriosis formulae (Shao, 1980; Cai, 1982; Lin *et al.*, 1988; Jin, 1991; Qu, 1992; Hu and Li, 1995; Cai *et al.*, 1999; Liu *et al.*, 2006; Yang *et al.*, 2006a). Curcuma zeodaria belongs to the plant family Zingiberaceae, which consists of ~80 different Curcuma species such as Curcuma longa (turmeric). Curcumin is a major active component of Curcuma. There are substantial *in vitro* and animal data indicating that curcumin has anti-inflammatory activity (Kumar *et al.*, 1998; Siddiqui *et al.*, 2006). Curcumin

was shown to suppress the NF-KappaB pathway and NF-KappaB target cytokine genes (Takada *et al.*, 2004) (Tables 1, 2, 5). Cao *et al.* demonstrated anti-inflammatory effects of curcumin in endometrial stromal cells; curcumin inhibited NF-Kappa B induction of a pro-inflammatory and angiogenic cytokine, the macrophage migration inhibitory factor (MIF) in this endometriosis *in vitro* model (Cao *et al.*, 2005). Recent studies could corroborate that curcumin arbitrates the effects by modulation of several important molecular targets, including NF-KappaB mediated gene expression (e.g. TNF, IL-1, IL-6), other transcription factors (e.g. AP-1, Egr-1, beta-catenin and PPAR-gamma), enzymes (e.g. COX2, iNOS), receptors (e.g. EGFR and HER2) and cell cycle proteins (e.g. cyclin D1 and p21) (Shishodia *et al.*, 2005). Substantial research has shown for other anti-endometriosis herbs (e.g. Chinese angelica) that they modulate cytokine secretion, including TNF-alpha as demonstrated in a variety of model systems (Tseng and Chang, 1992; Lee *et al.*, 1995; Jang *et al.*, 2001; Van Dien *et al.*, 2001; Xu *et al.*, 2002; Tipton *et al.*, 2003; Chen and Chen, 2004; Syrovets *et al.*, 2005; Liu *et al.*, 2006) (Table 5). Moreover, herbal combinations were shown to exert cytokine (e.g. TNF-alpha, IL-6, IL-8) suppressive effects studied in animal models of endometriosis (Yu *et al.*, 2000; Qu *et al.*, 2005). One example is YWN, a traditional anti-endometriosis formula (Table 3, 5), which decreased serum cytokine levels (e.g. TNF-alpha, IL-6 and IL-8) in a rodent model of endometriosis (Qu *et al.*, 2005).

A sophisticated new approach to create novel, low-toxicity anti-inflammatory drugs is to design chemical analogues of naturally occurring medicinal herbal compounds. For example, newly developed synthetic analogues of curcumin (e.g. EF24-tripeptide

Table 3: Clinical studies on herbal combinations used in the treatment of endometriosis

Herbal combination	Formula name	Patients (n)	Efficacy ^a	Language	Authors
Chinese angelica, corydalis, curcuma, persica, red peony, safflower, salvia root, tortoise shell	YWN versus gestrinone versus untreated	20 versus 19 versus 13	NS ^b	English	Yang <i>et al.</i> (2006a)
Zeodaria, salvia root versus danazol	Dan'e recipe versus danazol	189 versus 160	NS ^b	Chinese	Cai <i>et al.</i> (1999)
Curcuma, cuscuta, epimedium, eupolyphaga, leech, parganium, pangolin versus danazol	Neiyixiao Recipe versus danazol	58 versus 45	NS ^b	Chinese	Liu <i>et al.</i> (1998)
Persica, rhubarb, succinum, tortoise shell	Neiyi Recipe #2	48	<0.05	Chinese	Wang <i>et al.</i> (1998)
Curcuma, eupolyphaga, leech, liquidambar, pangolin, prunella, sappan, scales, sparganium		48	97.6	Chinese	Hu <i>et al.</i> (1995)
Cinnamon, poria		46	91.3%	Chinese	Liu <i>et al.</i> (1994)
Corydalis, curcuma, cyperus, loranthus, salvia, sparganium, trogopteris		54	94%	Chinese	Qu <i>et al.</i> (1992)
Rhubarb, succinum, tortoise shell	Neiyi Recipe #1	76	89%	Chinese	Wang <i>et al.</i> (1991)
Complex formula ^c		74	96%	Chinese	Li <i>et al.</i> (1991)
Complex formula ^d	–	45	84%	Chinese	Jin <i>et al.</i> (1990)
Calamus, Chinese angelica, gum, san-chi, trogopteris, typha	Shixiao Guijie Tang	30	93%	Chinese	Zhuang <i>et al.</i> (1990)
Complex formula ^e	Shofu Zhuyu Tang	40	97.5%	Chinese	Lin <i>et al.</i> (1988)
Complex formula ^f	Xiao Yan San	60	78%	Chinese	Liu <i>et al.</i> (1983)
Complex formulae ^g	I Xuefu Zhuyu Tang II Shaofu Zhuyu Tang III Taoren Chengqi Tang			English	Cao <i>et al.</i> (1983)
Complex formula ^h		43	88%	Chinese	Cai <i>et al.</i> (1982)
Complex formula ⁱ		30	80%	Chinese	Dai <i>et al.</i> (1982)
Complex formulae ^j		156	82%	Chinese	Shao (1980)

^aEfficacy rate (%) in terms of relief of dysmenorrhoeic symptoms.

^bNS, No significant difference between herbal combination and danazol.

^cCattail pollen, cinnamon, curculigo, dioscorea, epimedium, frankincense, ligustrum, lyceum, mielettia, myrrh, root bark, rehmannia and salvia aconite.

^dAngelica, artemisia anomala, asarum, cattail pollen, cinnamon twig, corydalis, Curcuma, gleditsia spine, hoelen, melia, moutan, persica, prunella, rubia, red peony, salvia, sparganium, trogopteris, myrrh and san-chi.

^eShaofu Zhuyu Tang: bulrush, Chinese angelica, cinnamon bark, cnidium, corydalis, fennel, dry ginger, myrrh, red peony and trogopteris.

^fXiao Yao San: achyranthes, artemisia, astragalus, bupleurum, Chinese angelica; cinnamon bark, cnidium, Curcuma, carbonized cyperus, frankincense, limestone, linden, myrrh, phellodendron, carbonized rhubarb, salvia, sanqui, sparganium, trogopteris and typha.

^gI: achyranthes bupleurum, carthamus, chih-ko, Chinese angelica, cnidium, licorice, persica, platycodon, raw rehmannia and red peony (additions: cattail pollen, Curcuma, sparganium, trogopteris); II: shaofu Zhuyu Tang: bulrush, Chinese angelica, cinnamon bark, cnidium, corydalis, dried ginger, fennel, myrrh, red peony, and trogopteris; III: taoren Chengqi Tang: achyranthes, Artemisia anomala, corydalis, moutan, persica, rhubarb, red peony, salvia.

^hChinese angelica, cinnamon, Curcuma, cyathula, cyperus, gleditsia spine, lacca, pangolin scales, salvia, sargassum, sparganium.

ⁱAgrimony, bupleurum, cattail pollen, corydalis, cyperus, melia, moutan, patrinia, prunella, oyster shell, red peony, raw rehmannia, salvia, saussurea, trogopteris.

^jGroup 1: bupleurum, bulrush, Chinese angelica, Curcuma, cyperus, gleditsia spine, pteropus, and sparganium, plus: centipede, earthworm, leech, tabanus; Group 2: salvia extract; Group 3: astragalus, bupleurum, cimicifuga bulrush, codonopsis, Curcuma, gleditsia spine, pteropus and sparganium; plus centipede, earthworm, leech, tabanus.

chloromethyl ketone) have been introduced to treat cancer and chronic inflammatory diseases (Selvam *et al.*, 2005; Otori *et al.*, 2006; Sun *et al.*, 2006). This alternative approach to target key cytokines may establish drugs that lack the deleterious side effects of conventional, as well as potential, endometriosis drugs such as

anti-TNF biologicals. Anti-TNF protein therapeutics were shown to have potentially severe side effects such as serious infection and to induce drug resistance in treated patients (Kenter and Cohen, 2006; Scott and Kingsley, 2006). Medicinal herbs or designed synthetic analogues of components of herbs with

Table 4: Toxicity and interactions of medicinal herbs used in the treatment of endometriosis (and typical clinical doses)

Herbs	Toxicity, Side effects	Interactions	LD ₅₀	Authors
Bupleurum 3–10 g	Low toxicity; abdominal distention, constipation, edema	Tolbutamide, interferon	1.19 g/kg in mice (i.p.)	Xie <i>et al.</i> (2006); Chen and Chen (2004); Ikegami <i>et al.</i> (2003)
Chinese angelica 5–15 g	Low toxicity; hypotension; bleeding; Stimulated MCF-7 cells	Warfarin	100 g/kg in mice	Hu <i>et al.</i> (2005); Amato <i>et al.</i> (2002); Page and Lawrence (1999); Huang (1998)
Cattail pollen (Typha) 3–10 g	Well-tolerated; very low toxicity; nausea, allergy vomiting, constipation		35.57 g/kg in mice (i.p.)	Chen and Chen (2004); Huang (1998)
Cinnamon twigs 1–2 g	Low toxicity		18.48 ± 1.8 g/kg in mice (i.v.)	Chen and Chen (2004); Perry <i>et al.</i> (1990); Hoskins (1984)
Cnidium fruit 3–9 g	Lox toxicity	Anticoagulants	65.86 g/kg (i.p.)	Li <i>et al.</i> (2001)
Corydalis 3–10 g	Decreases heart rate, lowers blood pressure; CNS depression	Barbiturates	100 ± 4.53 g/kg in mice	Chen and Chen (2004); Huang (1998)
Curcuma zedoaria 6–12 g	Relatively safe; nausea, vomiting, burning sensation, fatigue	Anticoagulants	16.75 g/kg in mice	Chen and Chen (2004); Huang (1998)
Cyperus 6–12 g	Malaise, sore throat, headache		1500 mg for cyperone in mice (i.p.)	Chen and Chen (2004)
Dahurian angelica 3–10 g	Increase of blood pressure; seizures, convulsions	Tolbutamide, diazepam, testosterone	53.82 g/kg in mice	Hu <i>et al.</i> (2005); Chen and Chen (2004); Ishihara <i>et al.</i> (2000)
Frankincense 3–10 g	Nausea, vomiting, epigastric pain			Ammon (2006); Chen and Chen (2004)
Licorice root 3–10 g	Hypertension, sodium retention, headache, hypokalemia	Digoxin, Corticosteroids	6.84 g/kg in mice (i.p.)	Zhou <i>et al.</i> (2004); Chen and Chen (2004); Elinav and Chajek-Shaul (2003)
Myrrh 3–10 g	Nausea vomiting			Chen and Chen (2004)
Persica 6–10 g	Cyanide poisoning	Anticoagulants	222 g/kg in mice	Chen and Chen (2004); Moing <i>et al.</i> (1987)
Poria 10–15 g	No significant side effects; diuretic	Diuretics	500 times the adult normal dose increased leucocytes (dogs)	Chen and Chen (2004)
Red peony root 6–15 g	Well-tolerated	Anticoagulants (warfarin)	Maximum safe dose is 50 g/kg in mice (i.v.)	Wong and Chan (2003)
Rhubarb 5–10 g	Nausea, vomiting, poor appetite, diarrhea;	Cardiac glycosides	153 g/kg in mice (oral)	Chen and Chen (2004); Wojcikowski <i>et al.</i> (2004b)
Salvia root 5–10 g	Relatively non-toxic	Anticoagulants, digoxin	80.5 ± 3.1 g/kg in mice	Hu <i>et al.</i> (2005); Chen and Chen (2004); Huang (1998)
Scutellaria 3–10 g	Hepatoprotective; diarrhea	Irrinotecan, beta-lactam antibiotics	3081 mg/kg of baicalein in mice (i.p.)	Hu <i>et al.</i> (2005); de Boer <i>et al.</i> (2005); Chen and Chen (2004); Takasuna <i>et al.</i> (1995)
Sparganium 3–10 g		Anticoagulants	233.9 ± 9.9 g/kg in mice (oral)	Chen and Chen (2004)
Tortoise shell 10–30 g	Allergy			Chen and Chen (2004)
Turmeric 3–10 g	Relatively safe, minimal toxicity; nausea, vomiting	Anticoagulants	50 times the adult dose for 30 days was not toxic in rats	Lao <i>et al.</i> (2006); Joshi <i>et al.</i> (2003); Chainani-Wu (2003)
White Peony root 5–10 g	Drowsiness, sedation	Anticoagulants, sedatives; antidiabetics	9.53 mg/kg in mice (i.p.)	Chen and Chen (2004)

cytokine-modulating effects may provide low-toxicity alternatives to treat endometriosis-associated symptoms (Adams *et al.*, 2004).

Effects of medicinal herbs, their active components and herbal combinations on the prostanoid pathway

The prostanoid pathway is suggested to be one of the key targets involved in the pathogenesis of endometriosis. Locally produced

PGE₂, a potent stimulator of aromatase, upregulates estrogen production, which in turn stimulates COX-2 to increase PGE₂ leading to increased endometrial cell proliferation in endometriosis (Noble *et al.*, 1997; Attar and Bulun, 2006). COX-2-selective inhibitors were shown to exert anti-proliferative effects (Hasegawa *et al.*, 2005), to decrease implant size in rodent models of endometriosis (Dogan *et al.*, 2004; Matsuzaki *et al.*, 2004) and to diminish endometriosis associated-pain in a clinical setting (Cobellis *et al.*,

Table 5: Cytokine suppressive effects of medicinal herbs and herbal formulae used in endometriosis

Herb/Combination	Major active component	Cytokines inhibited	Model	Author
White peony root	Paeoniflorin	TNF-alpha, IL-6	<i>In vitro</i>	Liu <i>et al.</i> (2006)
Frankincense	Boswellic acids	TNF-alpha	<i>In vitro</i>	Syrovets <i>et al.</i> (2005)
Myrrh		IL-6, IL-8	<i>In vitro</i>	Tipton <i>et al.</i> (2003)
Scutellaria	Wogonin	TNF-alpha	<i>In vitro</i>	Van Dien <i>et al.</i> (2001)
Curcuma	Curcumin	TNF-alpha	<i>In vitro</i>	Jang <i>et al.</i> (2001)
Poria		TNF-alpha, IL-1, IL-6	<i>In vitro</i>	Tseng <i>et al.</i> (1992)
YWN ^a	Formula	TNF-alpha, IL-6, IL-8	Murine	Qu <i>et al.</i> (2005)
Scutellaria	Wogonin	TNF-alpha	Murine	Van Dien <i>et al.</i> (2001)
Sparganium		IL-8	Murine	Lee <i>et al.</i> (1995)
Chinese angelica	Essential oils	TNF-alpha	Rabbit	Xu <i>et al.</i> (2002)

^aChinese angelica, corydalis, curcuma, persica, red peony, safflower, salvia root and tortoise shell.

2004). Among medicinal anti-endometriosis herbs that inhibit the prostanoid system, dahurian angelica root, cinnamon, licorice root, poria, scutellaria, Curcuma and the formulae KBG, YWN and Neiyi have been best studied (Wang *et al.*, 1991; Mori *et al.*, 1993; Yu *et al.*, 1993; Wang *et al.*, 1998; Giner-Larza *et al.*, 2000; Park *et al.*, 2001; Hong *et al.*, 2002; Huss *et al.*, 2002; Ban *et al.*, 2003; Prieto *et al.*, 2003; Furuhashi *et al.*, 2005; Qu *et al.*, 2006) (Table 6). Cinnamon and poria, both exhibiting COX-2 suppressive activity, are components of the anti-endometriosis herbal formula KBG (consisting of five different herbs cinnamon, poria, red peony, persica and tree peony bark). Treatment with the KBG formula suppressed spontaneous development of adenomyosis in a murine endometriosis model (Mori *et al.*, 1993). The beneficial effects of KBG in the murine endometriosis model may be attributed to the COX-2 inhibiting effect of KBG's constituents poria and cinnamon. YWN, an herbal composition (Chinese angelica, corydalis, curcuma, persica, red peony, safflower, salvia root, tortoise shell) (Table 6), reduced expression of COX-2 mRNA in endometriotic tissues in a rat model (Qu *et al.*, 2006). Furthermore, the anti-endometriosis herbal recipe Neiyi (Table 6) decreased prostaglandins (e.g. PGE₂, P_gF₁-alpha) levels in clinical settings in women with endometriosis (Wang *et al.*, 1991; Wang *et al.*, 1998; Wu *et al.*, 2000).

The highly regulated enzyme COX-2, which catalyzes prostaglandin production, has become a popular target for the development of new anti-inflammatory drugs. However, some of the COX-2 selective NSAIDs possess gastrointestinal side effects. In addition, recent data suggest that chronic use of COX-2 inhibitors is associated with increased cardiovascular risk (Vonkeman *et al.*, 2006). As a consequence, some of the COX-2 selective NSAIDs

were labelled with black box warning or withdrawn from the market. In the light of recent reports of serious unintended effects and black box warnings of COX-2 selective drugs, plants with direct or indirect suppressive effects on prostanoids (Huss *et al.*, 2002) represent a potential alternative resource of COX-2 selective inhibitors.

Effects of medicinal herbs, their active components and herbal combinations on oxidative status

Reactive oxygen species (ROS) are suggested to play a role in the pathogenesis of endometriosis (Murphy *et al.*, 1998). Retrograde menstruation allows transport of pro-oxidant factors, such as heme, iron and apoptotic endometrial cells, which are well-known inducers of oxidative stress, into the peritoneal cavity of women with endometriosis (Van Langendonck *et al.*, 2002). ROS can promote growth of endometrial stromal cells (Foyouzi *et al.*, 2004). Antioxidants, such as vitamin E, showed beneficial effects in an *in vitro* model of endometrial proliferation (Foyouzi *et al.*, 2004) and anti-inflammatory effects in a rodent model of LPS-induced inflammation leading to increased embryo viability (Mayorga *et al.*, 2004). Drugs with antioxidant properties, such as SPRMs, have been developed as possible treatment choices for endometriosis (Roberts *et al.*, 1996), but conclusive evidence on the benefits of the various modalities is lacking.

Commonly used 'anti-endometriosis' herbs such as Chinese angelica, curcuma and salvia root, turmeric and herbal formulae (e.g. KBG) have potent anti-oxidant effects (Cao *et al.*, 1996; Yoshioka *et al.*, 1998; Quiles *et al.*, 2002; hou *et al.*, 2004; Kang *et al.*, 2004; Moussaieff *et al.*, 2005; Sekiya *et al.*, 2005;

Table 6: Suppressive effects on the prostanoid pathway of medicinal herbs and combinations used for endometriosis

Herb/Combination	Major active component	Suppression of prostanoids	Model	Author
Licorice	Licochalcone A	Phospholipase-2, COX-2, PGE2	<i>In vitro</i>	Furuhashi <i>et al.</i> (2005)
Dahurian angelica	Coumarin, furucocoumarin	COX-2, PGE2	<i>In vitro</i>	Ban <i>et al.</i> (2003)
Curcuma	Curcumin, Ceta- (or ar-) turmerone	PGE2	<i>In vitro</i>	Hong <i>et al.</i> (2002)
Cinnamon	Cinnamonaldehyde	COX-2	<i>In vitro</i>	Huss <i>et al.</i> (2002)
YWN ^a	Formula	COX-2	Murine	Qu <i>et al.</i> (2006)
Poria	Pachymic acids	Phospholipase-A2	Murine	Prieto <i>et al.</i> (2003)
Scutellaria	Wogonin	COX-2	Murine	Park <i>et al.</i> (2001)
Neiyi ^b	Formula	PGE2, P _g F ₂ -alpha	Human	Wang <i>et al.</i> (1998)

^aChinese angelica, corydalis, curcuma, persica, red peony, safflower, salvia root and tortoise shell.

^bPersica, rhubarb, succinum and tortoise shell.

Zhou *et al.*, 2005; El-Ashmawy *et al.*, 2006) (Table 7). The ability of KBG and vitamin E to prevent atherosclerosis was compared in diet-induced hypercholesterolemic rabbits (Sekiya *et al.*, 2005). KBG had a stronger anti-oxidant effect than vitamin E shown in this animal model. The superiority of the herbal combination KBG to vitamin E in this animal likely relates to synergistic interactions within the herbal combination.

In this context, it is noteworthy to mention that although increased serum levels of vitamin E levels were associated with decreased risk for cancer and cardiovascular diseases (Stahelin *et al.*, 1991), exogenous oral administration of the antioxidant vitamin E did not show benefits in cancer and cardiovascular disease in human intervention studies (Brown *et al.*, 2001; Lichtenstein and Russell, 2005). Data from the HOPE and HOPE-TOO trials showed that vitamin E did not decrease fatal or non-fatal cancer (Lonn *et al.*, 2005). Vitamin E alone also did not reduce myocardial infarction, stroke, CVD death and other cardiovascular morbidities; instead vitamin E increased hospitalization for heart failure (Lonn *et al.*, 2005). In contrast to these findings, diets high in vitamin E showed beneficial influence on the incidence on cardiovascular health and cancer risk (Knoops *et al.*, 2004; Lichtenstein and Russell, 2005). In an elderly population (70–90 years), adherence to a Mediterranean diet and healthful lifestyle is associated with a more than 50% lower rate of all-causes and cause-specific mortality (HALE project) (Knoops *et al.*, 2004). In combination, these studies (HOPE and HALE projects) suggest that either vitamin E has limited or no benefit or that it has benefit only in combination with other food-related substances. In an Italian endometriosis population, higher intake of green vegetables and fresh fruit caused a significant reduction of the risk for endometriosis (Parazini *et al.*, 2004). Interpretation of nutritional studies on the effect of dietary patterns on cancer and cardiovascular disease prevention led to the definition of the ‘whole food’ concept (Temple and Galdwin, 2003). Several components in fruits and vegetables with beneficial effects are postulated to prevent disease via their synergistic effects. This ‘teamwork’ principle is likely to hold for synergistic herbal therapies as well, but research will be needed to delineate optimal combinations.

Anti-nociceptive of medicinal herbs, their active components and herbal combinations used in the treatment of endometriosis

The cellular and neural mechanisms of pelvic pain associated with endometriosis are poorly understood (Gambone *et al.*, 2002). It has been proposed that direct invasion of pelvic nerves by

endometriotic implants or nearby release of inflammatory humoral factors (e.g. prostaglandins, cytokines) stimulate sensory and parasympathetic afferents that carry the perceived painful stimuli (Tulandi *et al.*, 1998). Ectopic endometrial growths themselves develop autonomic and sensory innervations (Berkley *et al.*, 2005).

Drugs most commonly prescribed to treat endometriosis pain symptoms, particularly dysmenorrhea, are NSAIDs, combined oral contraceptives (COC), and other analgesics (e.g. paracetamol). Evidence of the efficacy of COC on dysmenorrhea remains to be determined (Proctor *et al.*, 2001), and there has been reluctance to use the COC at young ages because of possible long-term health risks. The evidence on the efficacy of NSAIDs in the treatment of endometriosis is inconclusive (Allen *et al.*, 2005). Moreover, therapy with NSAIDs can bear side effects. Women using NSAIDs and even over the counter painkillers such as acetaminophen need to be aware of the possibility that these drugs may cause serious unintended effects including increased risk of cardiovascular events, hypertension and gastric ulceration (Dedier *et al.*, 2002; Forman *et al.*, 2005; Johnsen *et al.*, 2005).

Since ancient times, herbal medicine has been used to relieve pain and discomfort from wounds and burns. Evidence confirms pain alleviating mechanisms of several medicinal herbs used for endometriosis (e.g. cnidium fruit, corydalis, curcuma, dahurian angelica, frankincense, myrrh and white peony root) and herbal formulae (e.g. Neiyi) (Tuttle *et al.*, 1989; Liu *et al.*, 1990; Yu *et al.*, 1993; Yu *et al.*, 1995; Dolara *et al.*, 1996; Wang *et al.*, 1998; Wei *et al.*, 1999; Zhang *et al.*, 2000; Tsai *et al.*, 2001; Navarro Dde *et al.*, 2002; Kimmatkar *et al.*, 2003; Tatsumi *et al.*, 2004; Yuan *et al.*, 2004; Zhou *et al.*, 2005) (Table 8).

Corydalis is one of the most commonly used herbs in pain related syndromes (Chen and Chen, 2004), and is also used in many Chinese formulae designed for the treatment of endometriosis-associated pain (Dai, 1982; Cao, 1983; Lin *et al.*, 1988; Jin, 1991; Qu, 1992; Wieser *et al.*, 2005a,b; Yang *et al.*, 2006a) (Table 3). Tetrahydropalmatine (THP) is thought to be the main active pharmacological component of corydalis (Brown *et al.*, 2001); THP has been shown to exhibit anti-inflammatory, sedative, analgesic, hypnotic and muscle relaxant properties shown *in vitro* and rodent models (Wei *et al.*, 1999; Chen and Chen, 2004).

Anti-nociceptive CAM treatments are gaining increased popularity for treatment of dysmenorrhoea in industrialized countries (Fugh-Berman and Kronenberg, 2003). Therapies shown to be potentially effective in the treatment of dysmenorrhoea include vitamin E (Butler and McKnight, 1955), vitamin B1 (Gokhale, 1996; Wilson and Murphy, 2001) and ω -3 fatty acids (fish oil) (Deutch, 1995). One randomized controlled trial was conducted

Table 7: Antioxidant effects of medicinal herbs and herbal combinations used in endometriosis

Herb	Major active component	Antioxidant effect	Model	Author
Chinese angelica	Essential oils	Increased of SOD ^a and catalase	<i>In vitro</i>	Hou <i>et al.</i> (2004)
Salvia root		Reduced ROS	<i>In vitro</i>	Kang <i>et al.</i> (2004)
Salvia root	Tanshinone II-A Lithospermic acid B	Reduced ROS	<i>In vitro</i>	Cao <i>et al.</i> (1996)
Myrrh		Increased glutathione S-transferase	Murine	El-Ashmawy <i>et al.</i> (2006)
KBG ^b	Formula	Reduced lipid peroxide levels	Rabbit	Sekiya <i>et al.</i> (2005)
Turmeric	Curcumin	Increased antioxidants	Rabbit	Quiles <i>et al.</i> (2002)

^aSOD, superoxide dismutase.

^bCinnamon, poria, red peony, persica, tree peony bark.

Table 8: Anti-nociceptive effects of medicinal herbs and combinations used in endometriosis

Herb	Major active component	Postulated mechanisms	Model	Author
Curcuma	Curcumenol		Murine	Navarro <i>et al.</i> (2002)
White peony root	Paeoniflorin	Opioid receptor (κ)	Murine	Tsai <i>et al.</i> (2001)
Corydalis	Tetrahydropalaminine	Opioid, dopaminergic, and GABAergic mechanisms	Murine	Wei <i>et al.</i> (1999)
Myrrh	Furanoeudesma-1,3-diene	Opioid mechanisms	Murine	Dolara <i>et al.</i> (1996)
Cnidium fruit	Tetramethylpyrazine	Blocked uterine responsiveness to PGE2	Murine	Tuttle <i>et al.</i> (1989)
Salvia root		CNS	Cat	Liu <i>et al.</i> (1990)
Corydalis	Tetrahydropalaminine	Opioid, dopaminergic, and GABAergic mechanisms	Human	Yuan <i>et al.</i> (2004)
Dahurian angelica		Opioid, dopaminergic, and GABAergic mechanisms	Human	Yuan <i>et al.</i> (2004)
Frankincense	Boswellia serrata		Human	Kimmatkar <i>et al.</i> (2003)
Neiyi ^a	Formula	Increased beta-endorphin levels	Human	Yu <i>et al.</i> (1995)
Neiyi ^a	Formula	Increased beta-endorphin levels	Human	Yu <i>et al.</i> (1993)

^aRhubarb, persica, succinum, tortoise shell (formulae are modified according syndromes).

to evaluate the efficacy of an herbal combination consisting of corydalis, Chinese angelica and white peony root on the efficacy of dysmenorrhoea (Kennedy *et al.*, 2006); however, the sample size of this trial was too small. Herbal therapy may serve as a potential alternative especially for young adults with endometriosis-associated pain syndromes.

Discussion

A significant increase in lifetime use of herbs and other natural products was noted from 12.1% to 18.6% between 1998 and 2002 in the USA (Tindle *et al.*, 2005). On 16 May 2002, the WHO released its first global strategy on traditional medicines and announced that it will be greatly extending its activities in non-allopathic therapies. Several important drugs are derived from natural products: aspirin (1889; from willow bark), penicillin (1940s; from fungus), artemisinin (1980; from quinghaosu), mevacor (1987; from fungus), taxol (1993; from the pacific yew tree) and byetta (2005; from Gila monster saliva). It is hoped that medicinal herbs or combinations of medicinal herbs and drugs will promote health and well being, while minimizing toxicities and side effects.

Alternative pain therapies have gained popularity particularly among women with endometriosis. It is predicted that CAM will be used in combination with allopathic medicine or will completely replace standard pharmacological options for some women with endometriosis as shown in an Australian population (Cox *et al.*, 2003). Despite the lack of rigorous evidence, many recipients of herbal therapy view these natural products to be safer and more effective alternatives to Western medicine. Other CAM users believe them to be more compatible with their beliefs even if they are less efficacious than conventional therapy (Astin *et al.*, 2000). In view of the need for alternatives in pain management, CAM therapies such as medicinal herbs and other botanicals with anti-inflammatory and pain-alleviating properties may be useful particularly in the treatment of endometriosis-associated symptoms. Herbal medicine has been prescribed safely by professionals in the USA for many years; however, there are only few formal studies on safety and efficacy of combining herbal combinations with prescription drugs.

Although clinical studies on herbs in the literature show promising effects, conclusive clinical evidence of the efficacy of medicinal herbs in the treatment of endometriosis-associated pain is

lacking. The effects of Chinese herbal medicine for treating endometriosis remain unclear for several reasons. First, most herbs currently used in human populations have been available for the last 2000 years and no efficient regulation mechanisms control their use (Wolsko *et al.*, 2005). Since 1994, phytomedicinal products (medicines derived from plants) are legally classified in the USA as dietary supplements. Included in this category are vitamins, minerals, herbs or other botanicals, amino acids and other dietary substances derived from animals or plants. The US Food and Drug Administration lacks quality standards (De Smet, 2005) and does not require levels of therapeutic evidence in the form of randomized clinical trials before such products are marketed to consumers. Some European countries, such as Germany and France, were among the first to introduce simplified registration procedures for herbal products. In order to harmonize different registration procedures, the European parliament has installed specific regulatory mechanisms covering medicinal herbal products (De Smet, 2005). Directive 2004/24/EC requires a special, but simplified registration procedure for certain medicinal products (De Smet, 2005). In addition, European legislation introduced a list of recognized herbs and mandates adverse-event reporting. Secondly, specific mechanisms of action of herbs are underreported and have not been tested using appropriate study designs including *in vitro* and animal models (Cao *et al.*, 2005; Qu *et al.*, 2005; Wieser *et al.*, 2005b). Finally, single herbs (e.g. Curcuma) and herbal combinations are composed of a number of bioactive compounds, complicating the investigation of their mechanisms of actions. Benefits of the inherent synergism within single herbs and herbal combinations are enhanced efficacy and reduced toxicity. Of course, liabilities are also possible as some interacting substances might compete with each other or increase toxicity. The potential for both positive and negative interactions necessitates careful study and validations of existing and new permutations.

Investigations of the efficacy, toxicity and herb-herb and drug-herb interactions should include testing in *in vitro* and rodent (e.g. rat) models and subhuman primate models (e.g. baboon) of endometriosis (Ryan *et al.*, 1994; D'Hooghe, 1997; Awwad *et al.*, 1999; Fazleabas *et al.*, 2002). Moreover, investigators should be encouraged to provide information on the purity, quality and composition of the herbs tested in experimental trials. Standardization would enhance interpretation of CAM data and thus the applicability of the findings. After acquiring mechanistic and safety

data, we propose that randomized clinical trials be planned and implemented using human subjects with endometriosis. Adolescents with endometriosis are the most likely to benefit from disease prevention, preservation of fertility and pain relief with innovative herbal treatments. The use of standard 'single drug' hormonal endometriosis regimens (e.g. GnRH-a, danazol and progestagens) is most likely to have untoward side effects and long-term morbidities in this important patient population.

Conclusion

Over the next 20 years, major shifts are likely to occur in the use of herbal compounds, vitamins, minerals, supplements, nutraceuticals or whole food nutrition. We should anticipate a wave of new natural medicines based on scientifically substantiated health claims. In the near future, we predict that more US adults will use medicinal herbs and herbal products to treat their diseases as we change our perspective from treating disease to promoting better health. Formal clinical trials testing the mechanisms of action, efficacy and toxicities of CAM therapies are needed. The establishment of the NCCAM Institute of the NIH was an important step towards the validation of popular and effective traditional treatments (Stokstad, 2000). Despite the limited growth in the NIH, the Fiscal Year (FY) 2006 President's budget for the NCCAM increased to \$122 692 000. NCCAM has sponsored several clinical trials on the efficacy of medicinal herbs in the treatment of irritable bowel syndrome, menopausal symptoms, hepatitis C, AIDS and cancer.

Controlled clinical studies will be needed to clarify the clinical efficacy of natural medicinal herbs or synthetic herbal analogues in the treatment of endometriosis-associated pain and investigate herb-herb and drug-herb interactions and other toxicities. A potential benefit of herbal therapy is the likelihood of synergistic interactions within individual (e.g. Curcuma) or combinations of plants and the constraint of untoward side effects. Experimental and epidemiological data (e.g. HOPE and HALE trials) infer that health promotion involves addressing many important determinants rather than adjusting the level of a single substance. We, therefore, believe that it is critical to establish streamlined registration procedures for natural medicinal products, particularly herbal ones that have a long tradition of safe use.

Acknowledgements

Supported by Erwin Schroedinger Auslandsstipendium (J2288-B13) and Emory University Research Committee (2007068).

References

Adams BK, Ferstl EM, Davis MC, Herold M, Kurtkaya S, Camalier RF, Hollingshead MG, Kaur G, Sausville EA, Rickles FR *et al.* Synthesis and biological evaluation of novel curcumin analogs as anti-cancer and anti-angiogenesis agents. *Bioorg Med Chem* 2004;**12**:3871–3883.

Allen C, Hopewell S, Prentice A, Allen C. Non-steroidal anti-inflammatory drugs for pain in women with endometriosis. *Cochrane Database Syst Rev* 2005;CD004753.

Amato P, Christophe S, Mellon PL. Estrogenic activity of herbs commonly used as remedies for menopausal symptoms. *Menopause* 2002;**9**:145–150.

Ammon HP. Boswellic acids in chronic inflammatory diseases. *Planta Med* 2006;**72**:1100–1116.

Amsterdam LL, Gentry W, Jobanputra S, Wolf M, Rubin SD, Bulun SE. Anastrozole and oral contraceptives: a novel treatment for endometriosis. *Fertil Steril* 2005;**84**:300–304.

Anderson FW, Johnson CT. Complementary and alternative medicine in obstetrics. *Int J Gynaecol Obstet* 2005;**91**:116–124.

Astin JA, Pelletier KR, Marie A, Haskell WL. Complementary and alternative medicine use among elderly persons: one-year analysis of a Blue Shield Medicare supplement. *J Gerontol A Biol Sci Med Sci* 2000;**55**:M4–M9.

Attar E, Bulun SE. Aromatase and other steroidogenic genes in endometriosis: translational aspects. *Hum Reprod Update* 2006;**12**:49–56.

Awwad JT, Sayegh RA, Tao XJ, Hassan T, Awwad ST, Isaacson K. The SCID mouse: an experimental model for endometriosis. *Hum Reprod* 1999;**14**:3107–3111.

Badawy SZ, Etman A, Cuenca V, Montante A, Kaufman L. Effect of interferon alpha-2b on endometrioma cells *in vitro*. *Obstet Gynecol* 2001;**98**:417–420.

Ban HS, Lim SS, Suzuki K, Jung SH, Lee S, Lee YS, Shin KH, Ohuchi K. Inhibitory effects of furanocoumarins isolated from the roots of *Angelica dahurica* on prostaglandin E2 production. *Planta Med* 2003;**69**:408–412.

Barrier BF, Bates GW, Leland MM, Leach DA, Robinson RD, Propst AM. Efficacy of anti-tumor necrosis factor therapy in the treatment of spontaneous endometriosis in baboons. *Fertil Steril* 2004;**81**(Suppl 1): 775–779.

Bergqvist A, Bruse C, Carlberg M, Carlstrom K. Interleukin 1beta, interleukin-6, and tumor necrosis factor-alpha in endometriotic tissue and in endometrium. *Fertil Steril* 2001;**75**:489–495.

Berkley KJ, Rapkin AJ, Papka RE. The pains of endometriosis. *Science* 2005;**308**:1587–1589.

Boucher A, Lemay A, Akoum A. Effect of hormonal agents on monocyte chemotactic protein-1 expression by endometrial epithelial cells of women with endometriosis. *Fertil Steril* 2000;**74**:969–975.

Brown BG, Zhao XQ, Chait A, Fisher LD, Cheung MC, Morse JS, Dowdy AA, Marino EK, Bolson EL, Alaupovic P *et al.* Simvastatin and niacin, antioxidant vitamins, or the combination for the prevention of coronary disease. *N Engl J Med* 2001;**345**:1583–1592.

Burnett MA, Antao V, Black A, Feldman K, Grenville A, Lea R, Lefebvre G, Pinsonneault O, Robert M. Prevalence of primary dysmenorrhea in Canada. *J Obstet Gynaecol Can* 2005;**27**:765–770.

Butler EB, McKnight E. Vitamin E in the treatment of primary dysmenorrhoea. *Lancet* 1955;**268**:844–847.

Cai L, Shu Y, Xie H. Clinical and experimental study on the treatment of endometriosis with dan'e mixture. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 1999;**19**:159–161.

Cai XS, Sun TW, Cai Z, Feng DX, Lu YD, Zhou CZ. 43 cases of endometriosis treated by traditional Chinese medicine. *Shanghai Zhong Za Zhi* 1982;**4**:12–13.

Cao EH, Liu XQ, Wang JJ, Xu NF. Effect of natural antioxidant tanshinone II-A on DNA damage by lipid peroxidation in liver cells. *Free Radic Biol Med* 1996;**20**:801–806.

Cao LX. Endometriosis as treated by traditional Chinese Medicine. *J Am Coll Tradit Chin Med* 1983;**1**:54–57.

Cao WG, Morin M, Metz C, Maheux R, Akoum A. Stimulation of macrophage migration inhibitory factor expression in endometrial stromal cells by interleukin 1, beta involving the nuclear transcription factor NFkappaB. *Biol Reprod* 2005;**73**:565–570.

Chainani-Wu N. Safety and anti-inflammatory activity of curcumin: a component of tumeric (*Curcuma longa*). *J Altern Complement Med* 2003;**9**:161–168.

Chen J, Chen T. *Chinese Medical Herbology and Pharmacology*. City of Industry, CA, USA: Art of Medicine Press; City of Industry, CA, USA, 2004.

Chwalisz K, Perez MC, Demanno D, Winkel C, Schubert G, Elger W. Selective progesterone receptor modulator development and use in the treatment of leiomyomata and endometriosis. *Endocr Rev* 2005;**26**:423–438.

Cobellis L, Razzi S, De Simone S, Sartini A, Fava A, Danero S, Gioffre W, Mazzini M, Petraglia F. The treatment with a COX-2 specific inhibitor is effective in the management of pain related to endometriosis. *Eur J Obstet Gynecol Reprod Biol* 2004;**116**:100–102.

Comar KM, Kirby DF. Herbal remedies in gastroenterology. *J Clin Gastroenterol* 2005;**39**:457–468.

Corson SL, Bolognese RJ. Ibuprofen therapy for dysmenorrhea. *J Reprod Med* 1978;**20**:246–252.

Cox H, Henderson L, Wood R, Cagliarini G. Learning to take charge: women's experiences of living with endometriosis. *Complement Ther Nurs Midwifery* 2003;**9**:62–68.

Crosignani P, Olive D, Bergqvist A, Luciano A. Advances in the management of endometriosis: an update for clinicians. *Hum Reprod Update* 2006;**12**:179–189.

D'Hooghe TM. Clinical relevance of the baboon as a model for the study of endometriosis. *Fertil Steril* 1997;**68**:613–625.

- Dai DY. 30 cases of endometriosis treated by taking Chiense herbs orally, externally, and by enema. *Shanghai Zhing Yi Yao Za Zhi* 1982;**3**:34–35.
- de Boer JG, Quiney B, Walter PB, Thomas C, Hodgson K, Murch SJ, Saxena PK. Protection against aflatoxin-B1-induced liver mutagenesis by *Scutellaria baicalensis*. *Mutat Res* 2005;**578**:15–22.
- De Smet PA. Herbal medicine in Europe—relaxing regulatory standards. *N Engl J Med* 2005;**352**:1176–1178.
- Dedier J, Stampfer MJ, Hankinson SE, Willett WC, Speizer FE, Curhan GC. Nonnarcotic analgesic use and the risk of hypertension in US women. *Hypertension* 2002;**40**:604–608, discussion 601–603.
- Deutch B. Menstrual pain in Danish women correlated with low n-3 polyunsaturated fatty acid intake. *Eur J Clin Nutr* 1995;**49**:508–516.
- Dogan E, Saygili U, Posaci C, Tuna B, Caliskan S, Altunyurt S, Saatli B. Regression of endometrial explants in rats treated with the cyclooxygenase-2 inhibitor rofecoxib. *Fertil Steril* 2004;**82**(Suppl 3):1115–1120.
- Dolara P, Luceri C, Ghelardini C, Monserrat C, Aioli S, Luceri F, Lodovici M, Menichetti S, Romanelli M N. Analgesic effects of myrrh. *Nature* 1996;**379**:29.
- Eisenberg DM, Davis RB, Ettner SL, Appel S, Wilkey S, Van Rompay M, Kessler RC. Trends in alternative medicine use in the United States, 1990–1997: results of a follow-up national survey. *JAMA* 1998;**280**:1569–1575.
- El-Ashmawy IM, Ashry KM, El-Nahas AF, Salama OM. Protection by turmeric and myrrh against liver oxidative damage and genotoxicity induced by lead acetate in mice. *Basic Clin Pharmacol Toxicol* 2006;**98**:32–37.
- Elinav E, Chajek-Shaul T. Licorice consumption causing severe hypokalemic paralysis. *Mayo Clin Proc* 2003;**78**:767–768.
- Fazleabas AT, Brudney A, Gurates B, Chai D, Bulun S. A modified baboon model for endometriosis. *Ann N Y Acad Sci* 2002;**955**:308–317, discussion 340–302, 396–406.
- Flaws B. *Endometriosis, Infertility & Traditional Chinese Medicine: A Laywomens's Guide*. Boulder, CO, USA: Blue Poppy Press, 1989.
- Forman JP, Stampfer MJ, Curhan GC. Non-narcotic analgesic dose and risk of incident hypertension in US women. *Hypertension* 2005;**46**:500–507.
- Foyouzi N, Berkkanoglu M, Arici A, Kwintkiewicz J, Izquierdo D, Duleba AJ. Effects of oxidants and antioxidants on proliferation of endometrial stromal cells. *Fertil Steril* 2004;**82**(Suppl 3):1019–1022.
- Friedlander RL. The treatment of endometriosis with Danazol. *J Reprod Med* 1973;**10**:197–199.
- Fugh-Berman A, Kronenberg F. Complementary and alternative medicine (CAM) in reproductive-age women: a review of randomized controlled trials. *Reprod Toxicol* 2003;**17**:137–152.
- Furuhashi I, Iwata S, Shibata S, Sato T, Inoue H. Inhibition by licochalcone A, a novel flavonoid isolated from liquorice root, of IL-1beta-induced PGE2 production in human skin fibroblasts. *J Pharm Pharmacol* 2005;**57**:1661–1666.
- Gambone JC, Mittman BS, Munro MG, Scialli AR, Winkel CA. Consensus statement for the management of chronic pelvic pain and endometriosis: proceedings of an expert-panel consensus process. *Fertil Steril* 2002;**78**:961–972.
- Giner-Larza EM, Manez S, Giner-Pons RM, Carmen Recio M, Rios JL. On the anti-inflammatory and anti-phospholipase A(2) activity of extracts from lanostane-rich species. *J Ethnopharmacol* 2000;**73**:61–69.
- Gokhale LB. Curative treatment of primary (spasmodic) dysmenorrhoea. *Indian J Med Res* 1996;**103**:227–231.
- Harrison TR, Fauci AS. *Harrison's Principles of Internal Medicine*, 14th edn. New York: McGraw-Hill, Health Professions Division New York, 1998.
- Hasegawa K, Ohashi Y, Ishikawa K, Yasue A, Kato R, Achiwa Y, Nishio E, Udagawa Y. Expression of cyclooxygenase-2 in uterine endometrial cancer and anti-tumor effects of a selective COX-2 inhibitor. *Int J Oncol* 2005;**26**:1419–1428.
- Hong CH, Noh MS, Lee WY, Lee SK. Inhibitory effects of natural sesquiterpenoids isolated from the rhizomes of *Curcuma zedoaria* on prostaglandin E2 and nitric oxide production. *Planta Med* 2002;**68**:545–547.
- Hoskins JA. The occurrence, metabolism and toxicity of cinnamic acid and related compounds. *J Appl Toxicol* 1984;**4**:283–292.
- Hou YZ, Zhao GR, Yang J, Yuan YJ, Zhu GG, Hiltunen R. Protective effect of Ligusticum chuanxiong and Angelica sinensis on endothelial cell damage induced by hydrogen peroxide. *Life Sci* 2004;**75**:1775–1786.
- Hu GZ, Li XY. 48 cases with endometriosis treated by the principle of eliminating stagnation and blood stasis and ecchymosis. *Shanghai Zhong Yi Yao Za Zhi* 1995;**2**:38–40.
- Hu Z, Yang X, Ho PC, Chan SY, Heng PW, Chan E, Duan W, Koh HL, Zhou S. Herb-drug interactions: a literature review. *Drugs* 2005;**65**:1239–1282.
- Huang K. *The Pharmacology of Chinese Herbs*, 2nd edn. USA: CRC Press, 1998.
- Hurxthal LM, Smith AT. Treatment of endometriosis and other gynecologic conditions with large doses of estrogens. *N Engl J Med* 1952;**247**:339–343.
- Huss U, Ringbom T, Perera P, Bohlin L, Vasange M. Screening of ubiquitous plant constituents for COX-2 inhibition with a scintillation proximity based assay. *J Nat Prod* 2002;**65**:1517–1521.
- Ikegami F, Fujii Y, Ishihara K, Satoh T. Toxicological aspects of Kampo medicines in clinical use. *Chem Biol Interact* 2003;**145**:235–250.
- Ishihara K, Kushida H, Yuzurihara M, Wakui Y, Yanagisawa T, Kamei H, Ohmori S, Kitada M. Interaction of drugs and Chinese herbs: pharmacokinetic changes of tolbutamide and diazepam caused by extract of Angelica dahurica. *J Pharm Pharmacol* 2000;**52**:1023–1029.
- Jang MK, Sohn DH, Ryu JH. A curcuminoid and sesquiterpenes as inhibitors of macrophage TNF-alpha release from *Curcuma zedoaria*. *Planta Med* 2001;**67**:550–552.
- Jin JL. 45 Cases of endometriosis treated by blood circulation promoting and stasis removing therapy and laboratory analysis for nail-fold microcirculation. *Shanxi Zhongyi Za Zhi* 1990;**11**:402–403.
- Johnsen SP, Larsson H, Tarone RE, McLaughlin JK, Norgard B, Friis S, Sorensen HT. Risk of hospitalization for myocardial infarction among users of rofecoxib, celecoxib, and other NSAIDs: a population-based case-control study. *Arch Intern Med* 2005;**165**:978–984.
- Joshi J, Ghaisas S, Vaidya A, Vaidya R, Kamat DV, Bhagwat AN, Bhide S. Early human safety study of turmeric oil (*Curcuma longa* oil) administered orally in healthy volunteers. *J Assoc Physicians India* 2003;**51**:1055–1060.
- Kang DG, Oh H, Sohn EJ, Hur TY, Lee KC, Kim KJ, Kim TY, Lee HS. Lithospermic acid B isolated from *Salvia miltiorrhiza* ameliorates ischemia/reperfusion-induced renal injury in rats. *Life Sci* 2004;**75**:1801–1816.
- Kelly JP, Kaufman DW, Kelley K, Rosenberg L, Anderson TE, Mitchell AA. Recent trends in use of herbal and other natural products. *Arch Intern Med* 2005;**165**:281–286.
- Kennedy S, Bergqvist A, Chapron C, D'Hooghe T, Dunselman G, Greb R, Hummelshoj L, Prentice A, Saridogan E. ESHRE guideline for the diagnosis and treatment of endometriosis. *Hum Reprod* 2005;**20**:2698–2704.
- Kennedy S, Jin X, Yu H, Zhong S, Magill P, van Vliet T, Kistemaker C, Voors C, Pasman W. Randomized controlled trial assessing a traditional Chinese medicine remedy in the treatment of primary dysmenorrhea. *Fertil Steril* 2006;**86**:762–764.
- Kenter MJ, Cohen AF. Establishing risk of human experimentation with drugs: lessons from TGN1412. *Lancet* 2006;**368**:1387–1391.
- Kimmatkar N, Thawani V, Hingorani L, Khiyani R. Efficacy and tolerability of Boswellia serrata extract in treatment of osteoarthritis of knee—a randomized double blind placebo controlled trial. *Phytomedicine* 2003;**10**:3–7.
- Kistner RW. The use of newer progestins in the treatment of endometriosis. *Am J Obstet Gynecol* 1958;**75**:264–278.
- Knoops KT, de Groot LC, Kromhout D, Perrin AE, Moreiras-Varela O, Menotti A, van Staveren WA. Mediterranean diet, lifestyle factors, and 10-year mortality in elderly European men and women: the HALE project. *JAMA* 2004;**292**:1433–1439.
- Kuboniwa S, Maemura S, Mimematsu M, Onishi K, Saigou F, Aruga F, Miwa Y. Mutagenicity studies of Kaishi-bukuryo-gan (TJ-25). *Jpn Pharmacol Ther* 1999;**27**:95–100.
- Kumar A, Dhawan S, Hardegen NJ, Aggarwal BB. Curcumin (Diferuloylmethane) inhibition of tumor necrosis factor (TNF)-mediated adhesion of monocytes to endothelial cells by suppression of cell surface expression of adhesion molecules and of nuclear factor-kappaB activation. *Biochem Pharmacol* 1998;**55**:775–783.
- Kuohung W, Jones GL, Vitonis AF, Cramer DW, Kennedy SH, Thomas D, Hornstein MD. Characteristics of patients with endometriosis in the United States and the United Kingdom. *Fertil Steril* 2002;**78**:767–772.
- Lao CD, Ruffin MT, Normolle D, Heath DD, Murray SI, Bailey JM, Boggs ME, Crowell J, Rock CL, Brenner De. Dose escalation of a curcuminoid formulation. *BMC Complement Altern Med* 2006;**6**:10.
- Lebovic DI, Mueller MD, Taylor RN. Immunobiology of endometriosis. *Fertil Steril* 2001;**75**:1–10.
- Lebovic DI, Kir M, Casey CL. Peroxisome proliferator-activated receptor-gamma induces regression of endometrial explants in a rat model of endometriosis. *Fertil Steril* 2004;**82**(Suppl 3):1008–1013.

- Lee GI, Ha JY, Min KR, Nakagawa H, Tsurufuji S, Chang IM, Kim Y. Inhibitory effects of Oriental herbal medicines on IL-8 induction in lipopolysaccharide-activated rat macrophages. *Planta Med* 1995;**61**:26–30.
- Lemay A, Quesnel G. Potential new treatment of endometriosis: reversible inhibition of pituitary-ovarian function by chronic intranasal administration of a luteinizing hormone-releasing hormone (LH-RH) agonist. *Fertil Steril* 1982;**38**:376–379.
- Li XY. Method of kidney tonifying and stasis removing for 74 cases of endometriosis. *Shanghai zhangyi za zhi* 1991;**7**:20–21.
- Li M, Handa S, Ikeda Y, Goto S. Specific inhibiting characteristics of tetramethylpyrazine, one of the active ingredients of the Chinese herbal medicine 'Chuanxiong,' on platelet thrombus formation under high shear rates. *Thromb Res* 2001;**104**:15–28.
- Lichtenstein AH, Russell RM. Essential nutrients: food or supplements? Where should the emphasis be? *JAMA* 2005;**294**:351–358.
- Lin YH, Cao HJ, Fu FZ, Yue GQ. An approach to treatment of endometriosis by traditional medicine. *Fujian Zhong yi Yao* 1988;**19**:21–23.
- Liu C, Shi W, Sun L, Zheng Q. Effects of radix *Salviae Miltiorrhizae* on visceral pain discharges in the posterior nucleus of the thalamus in cats. *Zhongguo Zhong Yao Za Zhi* 1990;**15**:112–115, 128.
- Liu JX. Clinical study of the treatment of endometriosis with traditional Chinese medicine. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 1994;**14**:337–339.
- Liu DF, Zhao HH, Shi YJ, Zheng ZM, Zhou T, Wang YQ. The treatment of endometriosis by "activating blood circulation". *Zhong Xi Yi Jie He Za Zhi* 1983;**3**:207–209.
- Liu J, Li X, Hu X. Clinical observations on treatment of endometriosis by tonifying kidney and removing blood stasis. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 1998;**18**:145–147.
- Liu DF, Wei W, Song LH. Protective effect of paeoniflorin on immunological liver injury induced by bacillus Calmette-Guerin plus lipopolysaccharide: modulation of tumour necrosis factor-alpha and interleukin-6 mRNA. *Clin Exp Pharmacol Physiol* 2006;**33**:332–339.
- Lonn E, Bosch J, Yusuf S, Sheridan P, Pogue J, Arnold JM, Ross C, Arnold A, Sleight P, Probstfield J *et al.* Effects of long-term vitamin E supplementation on cardiovascular events and cancer: a randomized controlled trial. *JAMA* 2005;**293**:1338–1347.
- Maas JW, Calhaz-Jorge C, ter Riet G, Dunselman GA, de Goeij AF, Struijker-Boudier HA. Tumor necrosis factor-alpha but not interleukin-1 beta or interleukin-8 concentrations correlate with angiogenic activity of peritoneal fluid from patients with minimal to mild endometriosis. *Fertil Steril* 2001;**75**:180–185.
- Maciocia G. *Obstetrics and Gynecology in Chinese Medicine*. New York, NY, USA: Churchill Livingstone, 1997.
- Matsuzaki S, Canis M, Darcha C, Dallel R, Okamura K, Mage G. Cyclooxygenase-2 selective inhibitor prevents implantation of eutopic endometrium to ectopic sites in rats. *Fertil Steril* 2004;**82**:1609–1615.
- Mayorga M, Iborra A, Estany S, Martinez PM. Protective effect of vitamin E in an animal model of LPS-induced inflammation. *Am J Reprod Immunol* 2004;**52**:356–361.
- Moing A, Salesses G, Saglio PH. Growth and the composition and transport of carbohydrate in compatible and incompatible peach/plum grafts. *Tree Physiol* 1987;**3**:345–354.
- Mori T, Sakamoto S, Singtripop T, Park MK, Kato T, Kawashima S, Nagasawa H. Suppression of spontaneous development of uterine adenomyosis by a Chinese herbal medicine, keishi-bukuryo-gan, in mice. *Planta Med* 1993;**59**:308–311.
- Mori T, Yamasaki S, Masui F, Matsuda M, Sasabe H, Zhou YF. Suppression of the development of experimentally induced uterine adenomyosis by a novel matrix metalloproteinase inhibitor, ONO-4817, in mice. *Exp Biol Med (Maywood)* 2001;**226**:429–433.
- Moussaieff A, Fride E, Amar Z, Lev E, Steinberg D, Gallily R, Mechoulam R. The Jerusalem Balsam: from the Franciscan Monastery in the old city of Jerusalem to Martindale 33. *J Ethnopharmacol* 2005;**101**:16–26.
- Murphy AA, Santanam N, Parthasarathy S. Endometriosis: a disease of oxidative stress? *Semin Reprod Endocrinol* 1998;**16**:263–273.
- Nap AW, Dunselman GA, Griffioen AW, Mayo KH, Evers JL, Groothuis PG. Angiostatic agents prevent the development of endometriosis-like lesions in the chicken chorioallantoic membrane. *Fertil Steril* 2005;**83**:793–795.
- Navarro Dde F, de Souza MM, Neto RA, Golin V, Niero R, Yunes RA, Delle Monache F, Cechinel Filho V. Phytochemical analysis and analgesic properties of *Curcuma zedoaria* grown in Brazil. *Phytomedicine* 2002;**9**:427–432.
- Noble LS, Takayama K, Zeitoun KM, Putman JM, Johns DA, Hinshelwood MM, Agarwal VR, Zhao Y, Carr BR, Bulen SE, I. Prostaglandin E2 stimulates aromatase expression in endometriosis-derived stromal cells. *J Clin Endocrinol Metab* 1997;**82**:600–606.
- Ohuri H, Yamakoshi H, Tomizawa M, Shibuya M, Kakudo Y, Takahashi A, Takahashi S, Kato S, Suzuki T, Ishioka C *et al.* Synthesis and biological analysis of new curcumin analogues bearing an enhanced potential for the medicinal treatment of cancer. *Mol Cancer Ther* 2006;**5**:2563–2571.
- Page RL, II, Lawrence JD. Potentiation of warfarin by dong quai. *Pharmacotherapy* 1999;**19**:870–876.
- Parazzini F, Chiaffarino F, Surface M, Chatenoud L, Cipriant S, Chiantera V, Benzi G, Fedele L. Selected food intake and risk of endometriosis. *Hum Reprod* 2004;**19**:1755–1799.
- Park BK, Heo MY, Park H, Kim HP. Inhibition of TPA-induced cyclooxygenase-2 expression and skin inflammation in mice by wogonin, a plant flavone from *Scutellaria radix*. *Eur J Pharmacol* 2001;**425**:153–157.
- Perry PA, Dean BS, Krenzelok EP. Cinnamon oil abuse by adolescents. *Vet Hum Toxicol* 1990;**32**:162–164.
- Prieto JM, Recio MC, Giner MC, Manez S, Giner-Larza EM, Rios JL. Influence of traditional Chinese anti-inflammatory medicinal plants on leukocyte and platelet functions. *J Pharm Pharmacol* 2003;**55**:1275–1282.
- Proctor ML, Roberts H, Farquhar CM. Combined oral contraceptive pill (OCP) as treatment for primary dysmenorrhoea. *Cochrane Database Syst Rev* 2001:CD002120.
- Qu JZ. Wen hua combined with ear-pressing therapy for the treatment of 54 endometriosis. *Shanxi Zhongyi Za Zhi* 1992;**1**:38–39.
- Qu F, Zhou J, Ma B. The effect of Chinese herbs on the cytokines of rats with endometriosis. *J Altern Complement Med* 2005;**11**:627–630.
- Qu F, Zhou J, Yang DX, Ma WG, Ma BZ. Effects of Yiweining Recipe on expressions of metalloproteinase-2 and cyclooxygenase-2 mRNAs in ectopic endometrium of rats with endometriosis. *Zhong Xi Yi Jie He Xue Bao* 2006;**4**:634–638.
- Quiles JL, Mesa MD, Ramirez-Tortosa CL, Aguilera CM, Battino M, Gil A, Ramirez-Tortosa MC. *Curcuma longa* extract supplementation reduces oxidative stress and attenuates aortic fatty streak development in rabbits. *Arterioscler Thromb Vasc Biol* 2002;**22**:1225–1231.
- Reese KA, Reddy S, Rock JA. Endometriosis in an adolescent population: the Emory experience. *J Pediatr Adolesc Gynecol* 1996;**9**:125–128.
- Roberts CP, Murphy AA, Santanam N, Parthasarathy S. The effect of RU 486 and related compounds on cultured macrophage differentiation and function. *Am J Obstet Gynecol* 1996;**175**:248–257, discussion 257–249.
- Ryan IP, Schriock ED, Taylor RN. Isolation, characterization, and comparison of human endometrial and endometriosis cells *in vitro*. *J Clin Endocrinol Metab* 1994;**78**:642–649.
- Scott DL, Kingsley GH. Tumor necrosis factor inhibitors for rheumatoid arthritis. *N Engl J Med* 2006;**355**:704–712.
- Sekiya N, Kainuma M, Hikiami H, Nakagawa T, Kouta K, Shibahara N, Shimada Y, Terasawa K. Oren-gedoku-to and Keishi-bukuryo-gan-ryo inhibit the progression of atherosclerosis in diet-induced hypercholesterolemic rabbits. *Biol Pharm Bull* 2005;**28**:294–298.
- Selak V, Farquhar C, Prentice A, Singla A. Danazol for pelvic pain associated with endometriosis. *Cochrane Database Syst Rev* 2001:CD000068.
- Selvam C, Jachak SM, Thilagavathi R, Chakraborti AK. Design, synthesis, biological evaluation and molecular docking of curcumin analogues as antioxidant, cyclooxygenase inhibitory and anti-inflammatory agents. *Bioorg Med Chem Lett* 2005;**15**:1793–1797.
- Shao GQ. Clinical and experimental research on 156 cases of endometriosis treated by therapy of promoting blood circulation and removing stasis. *Shanghai Zhong Yi Yao Za Zhi* 1980;**3**:4–6.
- Shishodia S, Sethi G, Aggarwal BB. Curcumin: getting back to the roots. *Ann N Y Acad Sci* 2005;**1056**:206–217.
- Siddiqui AM, Cui X, Wu R, Dong W, Zhou M, Hu M, Simms HH, Wang P. The anti-inflammatory effect of curcumin in an experimental model of sepsis is mediated by up-regulation of peroxisome proliferator-activated receptor-gamma. *Crit Care Med* 2006;**34**:1874–1882.
- Stahelin HB, Gey KF, Eichholzer M, Ludin E, Bernasconi F, Thurneysen J, Brubacher G. Plasma antioxidant vitamins and subsequent cancer mortality in the 12-year follow-up of the prospective Basel Study. *Am J Epidemiol* 1991;**133**:766–775.
- Stokstad E. Alternative medicine. Stephen Straus's impossible job. *Science* 2000;**288**:1568–1570.

- Sun A, Shoji M, Lu YJ, Liotta DC, Snyder JP. Synthesis of EF24-tripeptide chloromethyl ketone: a novel curcumin-related anticancer drug delivery system. *J Med Chem* 2006;**49**:3153–3158.
- Syrovetts T, Buchele B, Krauss C, Laumonier Y, Simmet T. Acetyl-boswellic acids inhibit lipopolysaccharide-mediated TNF- α induction in monocytes by direct interaction with I κ B kinases. *J Immunol* 2005;**174**:498–506.
- Takada Y, Bhardwaj A, Potdar P, Aggarwal BB. Nonsteroidal anti-inflammatory agents differ in their ability to suppress NF- κ B activation, inhibition of expression of cyclooxygenase-2 and cyclin D1, and abrogation of tumor cell proliferation. *Oncogene* 2004;**23**:9247–9258.
- Takasuna K, Kasai Y, Kitano Y, Mori K, Kobayashi R, Hagiwara T, Kakhata K, Hirohashi M, Nomura M, Nagai E *et al*. Protective effects of kampo medicines and baicalin against intestinal toxicity of a new anticancer camptothecin derivative, irinotecan hydrochloride (CPT-11), in rats. *Jpn J Cancer Res* 1995;**86**:978–984.
- Tatsumi S, Mabuchi T, Abe T, Xu L, Minami T, Ito S. Analgesic effect of extracts of Chinese medicinal herbs Moutan cortex and Coicis semen on neuropathic pain in mice. *Neurosci Lett* 2004;**370**:130–134.
- Temple NJ, Gladwin KK. Fruit, vegetables, and the prevention of cancer: research challenges. *Nutrition* 2003;**19**:467–470.
- Tindle HA, Davis RB, Phillips RS, Eisenberg DM. Trends in use of complementary and alternative medicine by US adults: 1997–2002. *Altern Ther Health Med* 2005;**11**:42–49.
- Tipton DA, Lyle B, Babich H, Dabbous M. *In vitro* cytotoxic and anti-inflammatory effects of myrrh oil on human gingival fibroblasts and epithelial cells. *Toxicol In Vitro* 2003;**17**:301–310.
- Tsai HY, Lin YT, Tsai CH, Chen YF. Effects of paeoniflorin on the formalin-induced nociceptive behaviour in mice. *J Ethnopharmacol* 2001;**75**:267–271.
- Tseng J, Chang JG. Suppression of tumor necrosis factor- α , interleukin-1 β , interleukin-6 and granulocyte-monocyte colony stimulating factor secretion from human monocytes by an extract of *Poria cocos*. *Zhonghua Min Guo Wei Sheng Wu Ji Mian Yi Xue Za Zhi* 1992;**25**:1–11.
- Tulandi T, Chen MF, Al-Took S, Watkin K. A study of nerve fibers and histopathology of postsurgical, postinfectious, and endometriosis-related adhesions. *Obstet Gynecol* 1998;**92**:766–768.
- Tuttle RS, Marmelstein L, Trad T, Reddy S, Radley T. *In vitro* uterine response to tetramethylpyrazine, the active constituent of chung chong (a traditional Chinese medicine). *Am J Obstet Gynecol* 1989;**161**:1319–1323.
- Van Dien M, Takahashi K, Mu MM, Koide N, Sugiyama T, Mori I, Yoshida T, Yokochi T. Protective effect of wogonin on endotoxin-induced lethal shock in D-galactosamine-sensitized mice. *Microbiol Immunol* 2001;**45**:751–756.
- Van Langendonck A, Casanas-Roux F, Donnez J. Oxidative stress and peritoneal endometriosis. *Fertil Steril* 2002;**77**:861–870.
- Vercellini P, Cortesi I, Crosignani PG. Progestins for symptomatic endometriosis: a critical analysis of the evidence. *Fertil Steril* 1997;**68**:393–401.
- Vonkeman HE, Brouwers JR, van de Laar MA. Understanding the NSAID related risk of vascular events. *BMJ* 2006;**332**:895–898.
- Wang DZ, Wang ZQ, Zhang ZF. Treatment of endometriosis with removing blood stasis and purgation method. *Zhong Xi Yi Jie He Za Zhi* 1991;**11**:524–526, 515.
- Wang D, Wang Z, Yu C. Endometriosis treated by the method of resolving blood stasis to eliminate obstruction in the lower-jiao. *J Tradit Chin Med* 1998;**18**:7–11.
- Wei F, Zou S, Young A, Dubner R, Ren K. Effects of four herbal extracts on adjuvant-induced inflammation and hyperalgesia in rats. *J Altern Complement Med* 1999;**5**:429–436.
- Wieser F, Burks-Wicks C, Vigne J, Cohen M, Gaeddert A, Taylor R. Extracts from a traditional Chinese herbal remedy inhibit cell proliferation and induce apoptosis in human endometrial and endometriosis cells. In: *Annual ASRM Meeting*, Quebec, Canada, 2005a.
- Wieser F, Burks-Wicks C, Vigne JL, Cohen M, Gaeddert A, Taylor RN. An herbal remedy inhibits the expression of the chemokine regulated upon activation, normal T cell expressed, and secreted in endometriotic cells. In: *World Conference of Endometriosis*, Maastricht, Netherlands, 2005b.
- Wieser F, Vigne JL, Ryan I, Hornung D, Djalali S, Taylor RN. Sulindac suppresses nuclear factor- κ B activation and RANTES gene and protein expression in endometrial stromal cells from women with endometriosis. *J Clin Endocrinol Metab* 2005c;**90**:6441–6447.
- Wilson ML, Murphy PA. Herbal and dietary therapies for primary and secondary dysmenorrhoea. *Cochrane Database Syst Rev* 2001:CD002124.
- Wojcikowski K, Johnson DW, Gobe G. Medicinal herbal extracts—renal friend or foe? Part two: herbal extracts with potential renal benefits. *Nephrology (Carlton)* 2004a;**9**:400–405.
- Wojcikowski K, Johnson DW, Gobe G. Medicinal herbal extracts—renal friend or foe? Part one: the toxicities of medicinal herbs. *Nephrology (Carlton)* 2004b;**9**:313–318.
- Wolsko PM, Solondz DK, Phillips RS, Schachter SC, Eisenberg DM. Lack of herbal supplement characterization in published randomized controlled trials. *Am J Med* 2005;**118**:1087–1093.
- Wong AL, Chan TY. Interaction between warfarin and the herbal product quilinggao. *Ann Pharmacother* 2003;**37**:836–838.
- Wu Y, Hua L, Jin Y. Clinical study on endometrial ovarian cyst treated by integrated laparoscopy and Chinese herbal medicine. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 2000;**20**:183–186.
- Xie Y, Lu W, Cao S, Jiang X, Yin M, Tang W. Preparation of bupleurum nasal spray and evaluation on its safety and efficacy. *Chem Pharm Bull (Tokyo)* 2006;**54**:48–53.
- Xu XY, Li LH, Wu LS, Zhao CL, Lin HY. Adjustment effect of Radix Astragalus and Radix Angelicae sinensis on TNF- α and bFGF on renal injury induced by ischemia reperfusion in rabbit. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 2002;**27**:771–773.
- Yang DX, Ma WG, Qu F, Ma BZ. Comparative study on the efficacy of Yiweining and Gestrinone for post-operational treatment of stage III endometriosis. *Chin J Integr Med* 2006a;**12**:218–220.
- Yang JH, Chen MJ, Wu MY, Chen YC, Yang YS, Ho HN. Decreased suppression of interleukin-6 after treatment with medroxyprogesterone acetate and danazol in endometrial stromal cells of women with adenomyosis. *Fertil Steril* 2006b;**86**:1459–1465.
- Yoshioka T, Fujii E, Endo M, Wada K, Tokunaga Y, Shiba N, Hohsho H, Shibuya H, Muraki T. Antiinflammatory potency of dehydrocurdione, a deoxyar-y-derived sesquiterpene. *Inflamm Res* 1998;**47**:476–481.
- Yu CQ, Wang DZ, Wang ZQ. Effect of endometriosis pill No. 2 on beta-endorphin and dynorphin in endometriosis. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 1993;**13**:3, 7–9.
- Yu CQ, Zhai MF, Yao RM. Effect of nei-yi recipe on plasma beta-endorphin levels during menstrual cycle in women with endometriosis. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 1995;**15**:6–8.
- Yu C, Liu Y, Peng X. IL-8 and IL-10 levels in endometriosis and regulated role of neiyifang on them. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 2000;**20**:603–605.
- Yuan CS, Mehendale SR, Wang CZ, Aung HH, Jiang T, Guan X, Shoyama Y. Effects of *Corydalis yanhusuo* and *Angelica dahurica* on cold pressor-induced pain in humans: a controlled trial. *J Clin Pharmacol* 2004;**44**:1323–1327.
- Zhang Y, Hou G, Yue Y. The analgesic action of semen coicis on severe functional dysmenorrhea—a sequential trial observation. *J Tradit Chin Med* 2000;**20**:293–296.
- Zhao D, Lebovic DI, Taylor RN. Long-term progestin treatment inhibits RANTES (regulated on activation, normal T cell expressed and secreted) gene expression in human endometrial stromal cells. *J Clin Endocrinol Metab* 2002;**87**:2514–2519.
- Zhou S, Koh HL, Gao Y, Gong ZY, Lee EJ. Herbal bioactivation: the good, the bad and the ugly. *Life Sci* 2004;**74**:935–968.
- Zhou L, Zuo Z, Chow MS. Danshen: an overview of its chemistry, pharmacology, pharmacokinetics, and clinical use. *J Clin Pharmacol* 2005;**45**:1345–1359.
- Zhuang B, Xia GC. Correlation between treatment of syndrome differentiation and basic body temperature 21 cases of endometriosis. *Shanxi Zhongyi Za Zhi* 1990;**11**:537.

Submitted on January 19, 2007; resubmitted on April 11, 2007; accepted on May 4, 2007